## UNIVERSITY OF CALIFORNIA, DAVIS

BERKELEY • DAVIS • IRVINE • LOS ANGELES • MERCED • RIVERSIDE • SAN DIEGO • SAN FRANCISCO

VETERINARY GENETICS LABORATORY SCHOOL OF VETERINARY MEDICINE ONE SHIELDS AVENUE DAVIS, CALIFORNIA 95616-8744





## PROGRESSIVE RETINAL ATROPHY (PRA) REPORT

SARA VEGA CAT46041 Case: TOBALABA 1815 31-May-2012 Date Received: 7510447 05-Jun-2012 Report Date: **CHILE** 6102-4926-0705-8040 Report ID: Verify report at https://www.vgl.ucdavis.edu/myvgl/verify.html Cat: BAYOFOXES CHRONOS OF PERBAST Reg: SBT 052611 022 DOB: **05/26/2011** Breed: SO Sex: M Microchip: 981020005217011 Sire: DAM Reg: SBT 100408 020 Dam' SIRE Reg: SBV 032207013

PRA-CEP290 Result	PRA-CRX Result
N/N	N/N

## Result Codes for PRA-CEP290

Normal, cat does not have rdAc mutation\*

N/dAc Carrier, cat has one copy of rdAc mutation. Breeding between carriers is expected to produce 25% affected

kittens

rdAc /rdAc Affected

rdAc is a progressive retinal atrophy that causes late-onset blindness.\* Affected cats are born with normal vision, show retinal degeneration at about 7 months and are blind by age 3-5 years. The condition is inherited as an autosomal recessive, and there is no treatment. \* This test only detects the mutation in the CEP290 gene known to cause PRA-rdAc in Abyssinian, Somali, Ocicat, Siamese and related breeds, American Curl, American Wirehair, Bengal, Cornish Rex, Munchkin, Singapura and Tonkinese.

## Result Codes for PRA-CRX

Normal, cat does not have Rdy mutation\*\*

NRdy Affected, cat has one copy of the Rdy mutation. This cat will produce affected kittens 50% of the time when bred

to a normal cat, or 75% of the time when bred to another cat with one copy of the Rdy mutation.

Rdy Rdy Affected, cat will always produce affected kittens.

Rdy is an early-onset retinopathy caused by a defective protein that is critical for eye development.\*\* Affected kittens display dilated pupils and sluggish pupillary reflexes around 2 weeks of age and often become blind by about 7 weeks of age. The condition is inherited as an autosomal dominant trait, and there is no treatment. \*\* This test only detects the mutation in the CRX gene known to cause PRA-Rdy in Abyssinian and Somali breeds.